

The EMBO Journal Vol 12, pp.973-986, 1993, Copyright © European Molecular Biology Organization

Targeting expression of keratinocyte growth factor to keratinocytes elicits striking changes in epithelial differentiation in transgenic mice

L Guo, QC Yu and E Fuchs

Howard Hughes Medical Institute, Department of Molecular Genetics and Cell Biology, University of Chicago, IL 60637.

Keratinocyte growth factor (KGF) is a member of the fibroblast growth factor (FGF) family. Synthesized by cells of the dermal component of skin, KGF's potent mitogenic activity is on the epidermal component, which harbors the receptors for this factor. To explore the possible role of KGF in mesenchymal-epithelial interactions in skin, we used a human keratin 14 promoter to target expression of human KGF cDNA to the stratified squamous epithelia of transgenic mice. Mice expressing KGF in their epidermis typically appeared frail and weak, and often had grossly wrinkled skin. These mice exhibited a gross increase in epidermal thickness accompanied by alterations in epidermal growth and differentiation. Most remarkably, animals displayed several striking and unexpected changes, including a marked suppression of hair follicle morphogenesis and suppression of adipogenesis. With age, some animals developed gross transformations in the tongue epithelium and in epidermis. In addition, they exhibited elevated salivation and their salivary glands showed signs of altered differentiation. Collectively, our findings provide new and important insights into the roles of KGF, implicating this potent growth factor in eliciting global effects not only on growth, but also on development and differentiation, of skin and other tissues. In particular, KGF seems to interfere with signalling of some mesenchymal-epithelial interactions.

- ▶ Similar articles found in:
[EMBO Journal Online](#)
[PubMed](#)
- ▶ [PubMed Citation](#)
- ▶ Search Medline for articles by:
[Guo, L](#) || [Fuchs, E](#)
- ▶ Alert me when:
[new articles cite this article](#)
- ▶ [Download to Citation Manager](#)

This article has been cited by other articles:

- Hines, M. D., Allen-Hoffmann, B. L. (1996). Keratinocyte Growth Factor Inhibits Cross-linked Envelope Formation and Nucleosomal Fragmentation in Cultured Human Keratinocytes. *J. Biol. Chem.* 271: 6245-6251 [[Abstract](#)] [[Full Text](#)]
- Oro, A. E., Higgins, K. M., Hu, Z., Bonifas, J. M., Epstein Jr., E. H., Scott, M. P. (1997). Basal Cell Carcinomas in Mice Overexpressing Sonic Hedgehog. *Science* 276: 817-821 [[Abstract](#)] [[Full Text](#)]
- Cook, P. W., Piepkorn, M., Clegg, C. H., Plowman, G. D., DeMay, J. M., Brown, J. R., Pittelkow, M. R. (1997). Transgenic Expression of the Human Amphiregulin Gene Induces a Psoriasis-like Phenotype. *J. Clin. Invest.* 100: 2286-2294 [[Abstract](#)] [[Full Text](#)]

- Bajaj-Elliott, M., Poulson, R., Pender, S. L.F., Wathen, N. C., MacDonald, T. T. (1998). Interactions between Stromal Cell- derived Keratinocyte Growth Factor and Epithelial Transforming Growth Factor in Immune-mediated Crypt Cell Hyperplasia. *J. Clin. Invest.* 102: 1473-1480 [[Abstract](#)] [[Full Text](#)]
- Charpentier, E., Lavker, R. M., Acquista, E., Cowin, P. (2000). Plakoglobin Suppresses Epithelial Proliferation and Hair Growth In Vivo. *J. Cell Biol.* 149: 503-520 [[Abstract](#)] [[Full Text](#)]
- (1999). Animal Models of Psoriasis - What Can We Learn from Them?. *J Invest Dermatol* 112: 405-410 [[Abstract](#)] [[Full Text](#)]
- Longan, L., Longnecker, R. (2000). Epstein-Barr virus latent membrane protein 2A has no growth-altering effects when expressed in differentiating epithelia. *J Gen Virol* 81: 2245-2252 [[Abstract](#)] [[Full Text](#)]
- Hom, Y. K., Young, P., Thomson, A. A., Cunha, G. R. (1998). Keratinocyte Growth Factor Injected into Female Mouse Neonates Stimulates Uterine and Vaginal Epithelial Growth. *Endocrinology* 139: 3772-3779 [[Abstract](#)] [[Full Text](#)]
- Reddig, P. J., Dreckschmidt, N. E., Zou, J., Bourguignon, S. E., Oberley, T. D., Verma, A. K. (2000). Transgenic Mice Overexpressing Protein Kinase C{ ϵ } in Their Epidermis Exhibit Reduced Papilloma Burden but Enhanced Carcinoma Formation after Tumor Promotion. *Cancer Res* 60: 595-602 [[Abstract](#)] [[Full Text](#)]
- Escalante-Alcalde, D., Recillas-Targa, F., Valencia, C., Santa-Olalla, J., Chávez, P., Marroquín, A., Gutiérrez-X, L., Gariglio, P., Covarrubias, L. (2000). Expression of E6 and E7 Papillomavirus Oncogenes in the Outer Root Sheath of Hair Follicles Extends the Growth Phase and Bypasses Resting at Telogen. *Cell Growth Differ* 11: 527-539 [[Abstract](#)] [[Full Text](#)]
- Celli, G., LaRochelle, W. J., Mackem, S., Sharp, R., Merlino, G. (1998). Soluble dominant-negative receptor uncovers essential roles for fibroblast growth factors in multi-organ induction and patterning. *EMBO J.* 17: 1642-1655 [[Abstract](#)] [[Full Text](#)]
- Suzuki, S., Ota, Y., Ozawa, K., Imamura, T. (2000). Dual-Mode Regulation of Hair Growth Cycle by Two Fgf-5 Gene Products. *J Invest Dermatol* 114: 456-463 [[Abstract](#)] [[Full Text](#)]
- Booth, C., Potten, C. S. (2000). Keratinocyte Growth Factor Increases Hair Follicle Survival Following Cytotoxic Insult. *J Invest Dermatol* 114: 667-673 [[Abstract](#)] [[Full Text](#)]
- Potten, C. S., O'Shea, J. A., Farrell, C. L., Rex, K., Booth, C. (2001). The Effects of Repeated Doses of Keratinocyte Growth Factor on Cell Proliferation in the Cellular Hierarchy of the Crypts of the Murine Small Intestine. *Cell Growth Differ* 12: 265-275 [[Abstract](#)] [[Full Text](#)]
- ANDREADIS, S. T., HAMOEN, K. E., YARMUSH, M. L., MORGAN, J. R. (2001). Keratinocyte growth factor induces hyperproliferation and delays differentiation in a skin equivalent model system. *FASEB J.* 15: 898-906 [[Abstract](#)] [[Full Text](#)]
- Krakowski, M. L., Kritzik, M. R., Jones, E. M., Krah, T., Lee, J., Arnush, M., Gu, D., Sarvetnick, N. (1999). Pancreatic Expression of Keratinocyte Growth Factor Leads to Differentiation of Islet Hepatocytes and Proliferation of Duct Cells. *Am J Pathol* 154: 683-691 [[Abstract](#)] [[Full Text](#)]
- Emoto, H., Tagashira, S., Mattei, M.-G., Yamasaki, M., Hashimoto, G., Katsumata, T., Negoro, T., Nakatsuka, M., Birnbaum, D., Coulier, F., Itoh, N. (1997). Structure and Expression of Human Fibroblast Growth Factor-10. *J. Biol. Chem.* 272: 23191-23194 [[Abstract](#)] [[Full Text](#)]
- Mehling, A., Loser, K., Varga, G., Metze, D., Luger, T. A., Schwarz, T., Grabbe, S., Beissert, S. (2001). Overexpression of CD40 Ligand in Murine Epidermis Results in Chronic Skin Inflammation and Systemic Autoimmunity. *J. Exp. Med.* 194: 615-628 [[Abstract](#)] [[Full Text](#)]